

## Energy Problems – Set 4

1

A satellite in a perfectly circular orbit is in *Uniform Circular Motion*.

- a) Using Newton's Second Law and the problem solving techniques from last unit, find an expression for the orbital velocity of a satellite a distance  $r$  from the center of the Earth.


Satellites in geosynchronous orbit always remain above the same geographic spot on the Earth's surface, making them extremely handy for communications.

- b) Calculate the radius of the geosynchronous orbit.  
c) Calculate the orbital velocity of the geosynchronous orbit.

The radius of the Earth is:  $R_{\text{Earth}} = 6.37 \times 10^6 \text{ m}$ .

The mass of the Earth is :  $M_{\text{Earth}} = 5.97 \times 10^{24} \text{ kg}$

The gravitaional constant:  $G = 6.67 \times 10^{-11} \text{ m}^3/(\text{kg s}^2)$



## Energy Problems – Set 4

2

The Zeronians live on a planet with a mass of  $5.0 \times 10^{23}$  kg, a radius of  $3.0 \times 10^6$  m. Their planet is rapidly running out of atmosphere (so there's no wind resistance) and, because they enjoy skydiving AND breathing, they have built a space-craft to leave.



a) Their space-craft, which weighs 10 kg (the zeronians are tiny), launched with an initial velocity of 3000 m/s. What will be its velocity at when it is  $4.0 \times 10^6$  m from the center of the planet?

b) Prove that the maximum altitude of a space craft launched with an initial velocity  $V_0$  from the surface of a planet is:

where  $R_p$  is the radius of the planet,  $M_p$  is the mass of the planet, and  $G$  is the gravitational constant.

c) If the ship launches with a low  $V_0$ , it will go up to  $H_{max}$ , stop and come back down. As  $V_0$  increases,  $H_{max}$  increases. Calculate the required initial velocity for the particle to go up and NOT come back down. (HINT: What is  $H_{max}$  if the particle never comes back?)

## Energy Problems – Set 4

3


Consider a satellite in a circular orbit of radius  $r$  about the Earth.

- Calculate the Total Mechanical Energy of the Satellite (Potential Energy plus Kinetic Energy)
- If the satellite is in low Earth orbit, 200 km above the Earth's surface, how much work must the satellite thrusters do to boost it to the Geosynchronous orbit?

The radius of the Earth is:  $R_{\text{Earth}} = 6.37 \times 10^6$  m.

The mass of the Earth is :  $M_{\text{Earth}} = 5.97 \times 10^{24}$  kg

The gravitaional constant:  $G = 6.67 \times 10^{-11}$  m<sup>3</sup>/(kg s<sup>2</sup>)




## Energy Problems – Set 4

4

Imagine a spacecraft traveling from the Earth (mass  $M_1$ ) to the Moon (mass  $M_2$ ). The spacecraft will feel the force of gravity from both objects. Close to the Earth, the Earth's gravitational force will be greater. Close to the Moon, the Moon's gravitational force will be greater. Somewhere in between is an equilibrium point where the force goes to zero.

- Use Newton's Universal Law of Gravitation to find the distance  $x$  where the net force goes to zero.
- The gravitational potential is the sum of the potentials from each body. Writing an expression for the potential at a distance  $x$  from  $M_1$ . Find the equilibrium point by locating any minima in the function.
- Sketch the potential function. Is the equilibrium point stable or unstable?



## Energy Problems – Set 4

While on its way to Jupiter in 1993, the Galileo spacecraft made a flyby of asteroid Ida. Images captured of Ida showed that the asteroid has a tiny moon of its own, since given the name Dactyl.

Ida's average Radius is about 19 km. Dactyl orbits Ida with an orbital period of 27h at an orbital radius of 95 km.

Assuming that both Ida and Dactyl are spherical, determine Ida's approximate mass and density.



## Energy Problems – Set 4

6

Suppose the Earth were suddenly to stop revolving around the Sun. The gravitational force would then pull it directly into the Sun. What would be the Earth's speed as it crashed (i.e., just when the Earth's *surface* hits the Sun's *surface*)? Assume the Earth starts at rest and the Sun doesn't move toward the earth.

$$M_{\text{sun}} = 1.99 \times 10^{30} \text{ kg}$$


$$R_{\text{sun}} = 695,500 \text{ km}$$

$$R_{\text{Earth}} = 6378.1 \text{ km}$$

$$M_{\text{Earth}} = 5.9742 \times 10^{24} \text{ kg}$$

## Energy Problems – Set 4

Determine the escape speed of a rocket from the far side of Ganymede, the largest of Jupiter's moons. The radius of Ganymede is  $2.640 \times 10^6$  m, and its mass is  $1.495 \times 10^{23}$  kg. The mass of Jupiter is  $1.900 \times 10^{27}$  kg, and the distance between the center of mass of Ganymede and the center of mass of Jupiter is  $1.071 \times 10^9$  m. Be sure to include the gravitational effect of Jupiter, but you can ignore the motion of Jupiter and Ganymede as they revolve about the systems center of mass.



## Energy Problems – Set 4

8

Two identical stars with mass  $M$  orbit around their center of mass. Each orbit is circular and has radius  $R$ , so that the two stars are always on opposite sides of the circle.

(a) Find the gravitational force of one star on the other.

(b) Find the orbital speed of each star and the period of the orbit.

(c) How much energy must be added to the system to separate the two stars to infinity, where they are at rest? Write your answer in terms of  $M$  and  $R$ . (Note: the addition of energy, however it physically happens, is tantamount to non-conservative work.)